

TRANSIT AND RADIAL VELOCITY SURVEY EFFICIENCY COMPARISON FOR A HABITABLE ZONE EARTH

CHRISTOPHER J. BURKE¹ & P. R. MCCULLOUGH^{2,3}

Submitted

ABSTRACT

Transit and radial velocity searches are two techniques for identifying nearby extrasolar planets to Earth that transit bright stars. Identifying a robust sample of these exoplanets around bright stars for detailed atmospheric characterization is a major observational undertaking. In this study we describe a framework that answers the question of whether a transit or radial velocity survey is more efficient at finding transiting exoplanets given the same amount of observing time. Within the framework we show that a transit survey’s window function can be approximated using the hypergeometric probability distribution. We estimate the observing time required for a transit survey to find a transiting Earth-sized exoplanet in the HZ with an emphasis on late type stars. We also estimate the radial velocity precision necessary to detect the equivalent HZ Earth-mass exoplanet that also transits when using an equal amount of observing time as the transit survey. We find that a radial velocity survey with $\sigma_{rv} \sim 0.6 \text{ m s}^{-1}$ precision has comparable efficiency in terms of observing time to a transit survey with the requisite photometric precision $\sigma_{phot} \sim 300 \text{ ppm}$ to find a transiting Earth-sized exoplanet in the HZ of late M dwarfs. For Super-Earths, a $\sigma_{rv} \sim 2.0 \text{ m s}^{-1}$ precision radial velocity survey has comparable efficiency to a transit survey with $\sigma_{phot} \sim 2300 \text{ ppm}$.

Subject headings: eclipses – methods:statistical – planetary systems – surveys – techniques:photometric – techniques:radial velocities

1. INTRODUCTION

Transiting extrasolar planets⁴ of bright host stars provide a unique opportunity for physical characterization in ways impossible for non-transiting planets that are also too close to their host stars for direct imaging. Currently, the transit and radial velocity methods dominate the growth in planet discoveries both overall and for planets suitable for detailed physical characterization⁵.

This paper addresses the question, “of these two methods, transits or radial velocities, which is more efficient at discovering planets that transit bright stars?” This is a complex question given the astrophysical effects (e.g. planet mass-radius relation, stellar mass-radius relation, stellar mass-luminosity relation, radial velocity jitter, stellar activity, etc.) and technical challenges (e.g. instrumentation, telescope aperture, available observing time, costs of operation, available funding opportunities, etc.) that influence the planet yield of a survey. To compare the performance of a radial velocity survey to that of a transit survey, we adopt a parametric model for each. With some simplifying assumptions, we quantify the precision of a radial velocity survey that will achieve the same level of search completeness as a transit survey using the same amount of observing time.

For any transit survey, the vast majority of observations of any particular star cannot contribute directly to

the detection of a transit, simply because they are made when the planet is not transiting. To be successful, a transit survey must make a sufficiently large number of independent observations in order that the probability $p_{\geq k}$ of witnessing k or more transit(s) matches or exceeds a design requirement. For example, the four-year Kepler mission was designed to witness $k \geq 3$ transits of Earth analogs with orbital periods $\lesssim 1$ year (Borucki et al. 2010). For a specific set of observation times of a transit survey, the *window function*, $p_{\geq k}(P)$, describes the visibility of the repetitive transit signal of any given period, i.e. the planet’s orbital period, P (Gaudi 2000). A survey’s window function can be computed numerically with high fidelity, either before or after the actual observations, by Monte Carlo simulations (Burke et al. 2006; von Braun et al. 2009).

However, analytic approximations to the window function are quicker and more convenient to compute than Monte Carlo simulations (Deeg et al. 2004; Beatty & Gaudi 2008; Fleming et al. 2008). In addition, an analytic approximation is readily inverted to predict the minimum observing time necessary to exceed a given probability $p_{\geq k}(P)$ of witnessing k or more transits of a given P . We show in the appendix that the hypergeometric distribution provides a high fidelity statistical approximation to the window function and its derivation gives physical insight into the design of a transit survey or interpretation of results from them. Most importantly we show how the transit survey window function is primarily dependent upon the number of independent observing epochs, which enables us to compare the performance of a transit survey to a radial velocity survey for detecting transiting planets.

For a radial velocity survey, the situation is quite different: nearly every measurement may contribute to the

Electronic address: christopher.j.burke@nasa.gov

¹ SETI Institute/NASA Ames Research Center, Moffett Field, CA 94035

² Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218

³ Department of Physics and Astronomy, Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218

⁴ Hereafter, “planets” means “extrasolar planets.”

⁵ <http://exoplanetarchive.ipac.caltech.edu>

planet detection because the star’s radial velocity can be measured at any time and is constantly changing in response to the orbiting planet. (We write “nearly” every measurement because multiple observations made at the same epoch could be redundant if one would suffice at that particular orbital phase.) To estimate the detection sensitivity of a radial velocity survey, we adopt the prescription outlined by Cumming (2004).

Our analysis ends at the phase of a survey when the significance of the detection reaches a suitable level such that follow-up observations to refine the ephemeris and confirm the detection are warranted. Thus, we neglect budgeting for the additional observational effort required to follow-up on a significant signal as the effort required depends very strongly on the nature of the signal and remaining ambiguities of the signal that must be ruled out. In the case of a transit survey a detection from a single to few events leads to ambiguity in the orbital period (Stevenson et al. 2012; Berta et al. 2013) that can be difficult to reconcile with low signal to noise detections. The numerous sources of astrophysical false positives to the transiting planet signal need to be investigated (Brown 2003; Torres et al. 2004; O’Donovan et al. 2006a) along with statistical validation (Torres et al. 2011; Fressin et al. 2011; Barclay et al. 2013) or mass measurement confirmation (Charbonneau et al. 2009; Queloz et al. 2009; Batalha et al. 2011). In the case of a radial velocity detection, in addition to the photometric confirmation of a transit signal (Henry et al. 2000; Charbonneau et al. 2000), typically additional radial velocity observations are required to refine the ephemeris to a suitable level for scheduling the photometric observations (Gillon et al. 2007; Kane et al. 2009; Winn et al. 2011). Also, radial velocity detections must contend with the astrophysical false positives induced by stellar variability (Saar & Donahue 1997; Boisse et al. 2011; Gomes da Silva et al. 2012).

We apply our prescription to address two representative case studies. First, we compare our models of the performance of a particular transiting-planet finder, *MEarth* and its southern extension, *MEarth-South* (Nutzman & Charbonneau 2008; Irwin et al. 2011), to that of a particular radial velocity instrument, *HARPS* (Bonfils et al. 2011) and its northern twin, *HARPS-N* (Cosentino et al. 2012; Covino et al. 2013), for detecting a Super-Earth transiting at the inner edge of the habitable zone (HZ) as a function of stellar mass. In the second case, we study the relative performances of these planet finding techniques for finding Earth-sized planets transiting at the inner edge of the HZ of M dwarf stars. In each case, we use the inner edge of the HZ simply because planets there will have shorter periods than any other planets in their respective HZs and hence will be more easily detected by either method.

This paper is organized as follows. In Section 2 the transit survey window function is introduced, and in the appendix we derive analytic approximations to it. This leads to simple approximations for the observing time required to reach a specified level of completeness of a transit survey. The detection sensitivity of a radial velocity survey is inverted in Section 3 in order to determine the observing time required for it to reach an appropriate level of completeness to compete with the transit survey. The implications of this study as applied to the detection

of a transiting Super-Earth and an Earth analog at the inner edge of the HZ in Section 4 and its conclusions are given in Section 5.

2. TRANSIT SURVEY PERFORMANCE

We can approach the statistical modeling of a transit survey window function in two ways. We can ask,

A) each time the planet transits, were we observing?

or

B) each time we observe, is the planet transiting?

For our statistical model, we want *independent* trials, so to form a trial, we group observations over an appropriate time interval. Previous work (e.g. Eq. A2 of Deeg et al. 2004) adopted approach A, where each independent planetary *orbital period* interval is a binomial trial with probability of our witnessing it equal to the average fraction of time spent observing f_{cov} . In approach B (see Appendix), the appropriate independent time interval is the *transit duration*,

$$\tau_{\text{dur}} = 0.058 \left(\frac{M_\star}{M_\odot} \right)^{-1/3} \left(\frac{R_\star}{R_\odot} \right) \left(\frac{P}{1 \text{ day}} \right)^{1/3} [\text{day}], \quad (1)$$

for arbitrary stellar mass, $M_\star$, radius, $R_\star$, and orbital period, P . We have assumed the average relative chord length, $\pi/4$ and eccentricity $e = 0$ (Gilliland et al. 2000; Seager & Mallén-Ornelas 2003). Effects of non-zero eccentricity on transit survey yield are addressed by Barnes (2007) and Burke (2008). Approach B has the advantage of being readily related to individual observations parceled into lengths of the transit duration. The window function is determined by the number of unique observing epochs, n , of duration τ_{dur} . A set of observation exposures at a cadence shorter than the transit duration will improve the signal to noise ratio of the detection, but count as a single parcel increment to n for the purposes of the window function.⁶ Thus, n is not necessarily the number of exposures (see Section 6.3).

Regardless of approach (A or B), the physical situation is the same: in the time span, T_{obs} , between the first and last observations of a particular star with a transiting planet of orbital period, P , the expected number of transits is⁷

$$M = T_{\text{obs}}/P. \quad (2)$$

From the probability, p_k , of witnessing exactly k transits (derived in the appendix), the probability, $p_{\geq k}$, of witnessing k or more transits is given by

$$p_{\geq k} = 1 - \sum_{i=0}^{k-1} p_i. \quad (3)$$

⁶ We are explicitly separating out the dependence of transit detection on the window function and signal to noise ratio (Gaudi 2000).

⁷ There are either $M = \lfloor T_{\text{obs}}/P \rfloor$ or $M = \lfloor T_{\text{obs}}/P \rfloor + 1$ transits where $\lfloor \cdot \rfloor$ is the truncate to nearest integer operation. For uniform and random relative phase of the planetary transits with respect to the set of observation epochs, the expectation value of $\langle M \rangle = T_{\text{obs}}/P$

A transit survey may require detection of $k \geq 2$ transits in order to know the period and hence predict future transits, or it may require $k \geq 3$ transits in order to validate the detection, i.e. confirm that the times separating the transits are all consistent with a particular orbital period.

In the appendix we have isolated the mathematics of approximating the window function analytically. Figure 1 shows the window function for a hypothetical transit survey that operates in a mode dedicated to a single field for 30 consecutive days with observations for 8 hours each night. The specifications for the hypothetical transit survey are the same as presented in Figure 1 of Beatty & Gaudi (2008) to demonstrate their analytical representation of a transit survey and is typical for the first generation of ground-based Hot Jupiter transit surveys (*HAT* – Bakos et al. (2007); *KELT* – Beatty et al. (2012); *OGLE* – Konacki et al. (2003); *Qatar* – Alsubai et al. (2011); *TrES* – Alonso et al. (2004); *WASP* – Collier Cameron et al. (2007); *XO* – McCullough et al. (2006); and others).

The window function presented in this study and others (e.g. Deeg et al. 2004; Beatty & Gaudi 2008) provide convenient approximations to the average performance of a transit survey. These approximations may be useful for design purposes and for real time updates to a transit survey in progress. Also, this study articulates that transit survey completeness, in terms of the window function, derives primarily from the number of distinct observing epochs, n , and only secondarily from the time span T_{obs} (see Figure 2 and discussion in Section 6.5)⁸. That is, observations can be spread over an observing season or years rather than concentrated in a single observing run. A consequence is that a longitudinally-distributed network (e.g. HAT-Net) is not necessarily more observationally efficient at finding planets per unit of observing time than a single site (e.g. WASP), all other factors being equal. However, a longitudinally-distributed network of N telescopes can retire a particular field of view of targets in a smaller number of days than could be accomplished by the same N telescopes located at a single site at the same latitude as the telescopes of the network. Also, continuous observations avoid the ambiguities (e.g. of orbital period) inherent in interpreting observations with many gaps where transits can hide.

Figure 3 shows the window function for a transit survey that operates in a mode cycling between targets such as employed by the *MEarth* project (Nutzman & Charbonneau 2008; Berta et al. 2013). *MEarth* surveys the brightest M dwarfs by cycling between targets to obtain several observation per target per night. The observation span is an observing season for each target. We simulate this mode of operation through specifying a time series with $n=200$ observations that is expected for a target from a single observatory over the course of an observing season including losses due to weather. As a proxy for an observing time series we begin with a continuous PWV time series downloaded from the NOAA Earth System Research Laboratory Ground-Based GPS Meteorology program⁹ (Wolfe & Gutman 2000) for 9 months starting Septem-




FIG. 1.— Left: exact numerical window function for a hypothetical ground-based transit survey that observes for 30 consecutive days with observations for 8 hours each night and requires two transits for detection (dashed black line). Average of the numerical window function using a 1-day moving average (solid black line) compared to the hypergeometric model (solid red line) from this study along with its Poisson model limit (dash blue line). The window function approximation of Beatty & Gaudi (2008) (solid blue line) is also shown. Right: same as left panel, but requiring three transits for detection.

ber, 1 2008 with a 30 minute cadence from Flagstaff, AZ. We use the celestial coordinates RA 7h Dec +30, optimized for the middle of the observing season and observatory latitude, to find the airmass for each PWV observations. Only values when the airmass < 1.8 , PWV < 10 mm (to crudely simulate when the weather conditions would be too poor for observations), and Sun altitude < -5 deg are used to simulate times when observations could be made. The window function assumes stellar parameters appropriate for a mid M dwarf star ($M_{\odot}=0.2 M_{\star}$ [$M_{\odot}$]; $R_{\star}=0.26 [R_{\odot}]$).

In addition to the window functions assuming at least $k \geq 1, 2, 3$, and 5 transit events are observed (solid lines from top to bottom, respectively), Figure 3 also shows the window function (solid orange line) which corresponds to the triggering mode of operation adopted by the *MEarth* project (Nutzman & Charbonneau 2008).

⁸ In general, $n \leq$ the number of exposures (see Section 6.3)

⁹ <http://www.gpsmet.noaa.gov>




FIG. 2.— Demonstrating the strong dependence of a transit survey window function on n compared to the weak dependence on T_{obs} (see Section 6.5). The following three observing scenarios have the same n , but vary in T_{obs} . We define the baseline window function for a hypothetical ground-based transit survey that observes for 7 hours each night for $T_{obs}=30$ days using the hypergeometric model (solid black line) and 1-day moving average of exact numerical model (dashed black line). Alternatively, we simulate a window function for a longitudinally-distributed observing network with half, $T_{obs}=15$ day, but increased observing coverage to equal n of the baseline case (hypergeometric model, solid red line, and moving-average numerical model, dashed red line). Finally, we simulate a seasonally-distributed transit survey window function with two observing campaigns of 15 consecutive observing days for 7 hours each night with the two campaigns separated by a year (hypergeometric model, solid blue line, and moving-average numerical model, dashed blue line). We compare the previous scenarios to a scenario with $2n$ from the baseline case using $T_{obs}=30$ day, but increased observing coverage each day (hypergeometric mode, solid yellow line, and moving-average numerical model, dashed yellow line). Showing results requiring $k \geq 2$ (left panel) and $k \geq 3$ (right panel) transits.

Real-time analysis of the data identifies a potential flux decrement for a target, and the observing scheduler interrupts cycling through targets and enters a high cadence follow-up and stare mode in order to get a high sensitivity detection of the transit egress. This has the potential of detecting a planet with a single transit and relaxed photometric precision, which extends the detectability to longer orbital periods and enables sampling more targets.




FIG. 3.— Exact numerical window function with $n=200$ requiring observing at least one, two, three, and five transits (solid black lines top to bottom) compared to the hypergeometric model (solid red lines) and its Poisson approximation model (dash blue lines) for a typical observing time series over the course of an observing season. The analytic window function employing the triggering on a single transit mode of operation (orange line) is also shown.

However, in practice for M dwarfs, the transit duration $\lesssim 1$ hr and the transit must be observed during the first portion of the transit event to allow sufficient time to commence the high cadence staring mode. Thus, τ_{dur} must be reduced to account for the triggering mode delay. Throughout, we reduce τ_{dur} by $\tau_{trig}=30$ min when simulating the triggering mode. For the mid M dwarf example shown in Figure 3, the triggering mode window function is superior to the $k = 2$ window function for $P > 4$ day.

3. RADIAL VELOCITY PERFORMANCE

In this section we summarize our metrics for the performance of a radial velocity survey. We follow the procedure by Cumming (2004) to determine the limiting semi-amplitude threshold, K , of a radial velocity survey for a given n observations, each of which has radial velocity precision, σ_{rv} . We use the performance of the Lomb-Scargle periodogram for detecting planetary orbits. For a given completeness probability, p_c , we invert Equation 24 of Cumming (2004) to determine the argument of the error function,

$$\Gamma = \text{erf}^{-1}(2p_c - 1). \quad (4)$$

The error function argument, Γ , relates the expectation of the periodogram in the presence of a signal,

$$\langle z_s \rangle = (\nu/2)(K^2/2\sigma_{rv}^2), \quad (5)$$

where $\nu = n-5$ is the degrees of freedom, to the detection threshold of the experiment,

$$z_d = (\nu/2)((m/F)^{2/\nu} - 1), \quad (6)$$

where m is the number of independent frequencies searched and F is the false alarm probability of the experiment. The relationship between Equations 4, 5, and 6 is given by

$$\langle z_s \rangle - z_d = 2\Gamma\sqrt{\langle z_s \rangle}. \quad (7)$$

Equations 4-7 are taken directly from Cumming (2004). We numerically solve Equation 7 to derive the limiting K/σ_{rv} that a radial velocity achieves for a given p_c and n . We adopt parameters following Cumming (2004) that the radial velocity survey has a $F = 0.001$ and that $m \approx n$ (Horne & Baliunas 1986). An additional assumption for the validity of these results is that the baseline of observations is longer than the orbital period of the planet. In this study we ignore the impact of uncorrected stellar activity for setting a noise floor on radial velocity precision (Saar & Donahue 1997; Boisse et al. 2011; Gomes da Silva et al. 2012), and the additional observations necessary to refine the orbital parameters with sufficient precision to make an accurate prediction for the epoch of transit (Kane et al. 2009). When analyzing the identification of transiting planets from a radial velocity detected sample of planets in this paper, we assume a purely geometric prior transit probability rather than the potentially 10-25% higher posterior transit probability for planets first detected with radial velocities (Stevens & Gaudi 2013).

4. DISCUSSION

With the results from the previous sections, we can evaluate a transit survey designed to detect a transiting planet and quantify the equivalent radial velocity survey that is expected to have the same detection capability for the same total amount of observing time. With the framework described in this paper, we assume both the radial velocity survey and the transit survey achieve n observations. We then determine which method using n (independent of how long it takes to achieve n observations) achieves a higher survey completeness level for the planet of interest. One opportunity provided by *JWST* will be an unprecedented capability for measuring the atmospheric constituents of potentially habitable planets (Valenti et al. 2005; Deming et al. 2009). Taking advantage of the *JWST* opportunity requires identifying the most viable transiting planets for follow up by pushing the limits of photometric and radial velocity precision. We first compare the performances of the *MEarth* transiting survey (Nutzman & Charbonneau 2008; Berta et al. 2013) and the *HARPS* M dwarf radial velocity survey (Bonfils et al. 2011) for detection of Super-Earth planets orbiting the brightest M dwarfs. Then we show general results for the design of future surveys pushing the detection limits down to Earth-sized planets. The steps for the comparison begin with determining the n necessary for a transit survey to reach a given percent completeness for an inner HZ planet. We then calculate the radial velocity precision required to detect the same planet with the same amount of observing time.

4.1. Bright M Dwarfs Hosting A Transiting HZ Super-Earth

As an example we calculate the performance for a survey modeled after the *MEarth* transit survey (Nutzman & Charbonneau 2008; Berta et al. 2011). The *MEarth* survey is focused on identifying transiting Super-Earths orbiting the brightest M dwarfs employing the triggering mode of detection along with ex post facto period-folding search for multiple transit events. Since *MEarth* is observing the brightest M dwarfs, which are

far apart on the sky, it typically can observe only a single target per telescope pointing. In order to specify n required for a transit survey, we outline the properties of the planets and their stellar hosts in Table 1. For a given stellar mass, M_* , the stellar radius, R_* , stellar surface gravity $\log g$, and stellar effective temperature, T_{eff} , are estimated by a spline fit to the calibration of MK spectral types Table 15.7 and 15.8 of Cox (2000). For the given stellar parameters, we determine the orbital period corresponding to the inner edge of the HZ, P_{HZ} , from the results of Kasting et al. (1993). Empirically for M dwarfs, we find that an equilibrium temperature T_{eq} equal to 273 K occurs at the inner edge of the Kasting et al. (1993) HZ for a planetary bond albedo, $a = 0.3$, and full redistribution of the stellar flux, $f = 1$. Although a warmer $T_{\text{eq}} = 288$ K is a better match to the inner HZ edge of Kasting et al. (1993) for higher mass stars, $M_* > 0.4 M_{\odot}$, we use the cooler $T_{\text{eq}} = 273$ K to define the inner HZ for all stellar types throughout this work.

With the stellar parameters and P_{HZ} given, the additional quantities follow (τ_{dur} , transit probability, p_{tr} , radial velocity semi-amplitude, K , and transit depth, Δ) for an Earth-mass and Earth-radius analog as well as a Super-Earth mass and radius analog to GJ 1214 (Charbonneau et al. 2009; Berta et al. 2011). Table 1 shows n necessary for a transit survey to achieve a completeness probability $p_c = 0.75$ of witnessing a transit at least once in the triggering mode of operation at P_{HZ} . We assume $T_{\text{obs}}=200$ day or $T_{\text{obs}}=8 \times P$ whichever is greater. Focusing on the $0.2 M_*$ star, $n = 449$ are required with sufficient precision to detect the corresponding transit depth, $\Delta_{\text{Super-}\oplus} = 9233$ ppm. If every star has a planet orbiting with a period of approximately, P_{HZ} , then with $p_{\text{tr}} = 0.0179$, $1/p_{\text{tr}} = 56$ such targets need to be observed. At least $n/(p_{\text{tr}}p_c) \sim 34000$ observations are required to expect to find one transiting planet at the inner HZ orbiting a $0.2 M_*$ M dwarf. Cycling between targets with a cadence of 5 min and 8 hours of observations, reaching 34000 observations requires 355 clear nights. The above results are first-order guidance assuming unit planet occurrence. They can be conveniently scaled to the measured occurrence rates of planets orbiting M-dwarfs (Bonfils et al. 2011; Dressing & Charbonneau 2013; Berta et al. 2013). Fortunately, the targets are independent, so this amount of observing time can be achieved with multiple telescopes working in parallel (*MEarth* employs eight telescopes and is commissioning eight more in the Southern hemisphere) or observing fainter stars with multiple targets per field. Of course, if the fraction of stars with such a planet is less than unity, the required observing time will be more.

It is interesting to note that n is not monotonically increasing with M_* because the increasing HZ period is offset by the increase in τ_{dur} with increasing M_* . In addition, the trigger mode delay becomes an appreciable fraction of τ_{dur} for the lowest mass stars. For $M_* < 0.6 M_{\odot}$, $n \sim 500$, thus in principle similar number of observations are needed to find a HZ planet orbiting a late K as late M star in the triggering mode of operation. However, the minimum number of clear nights necessary to achieve the requisite number of observations $n_{\text{ngt}} = \tau_{\text{dur}}n/\tau_{\text{ngt}}$, assuming $\tau_{\text{ngt}} = 8$ hr observing each night is a monotonically increasing function of M_* . In addition,

the depth of the transit for fixed planet size is ~ 6 times smaller for late K hosts than late M hosts placing more stringent requirements on the photometric precision for observations of late K hosts.

For an individual target we determine the radial velocity precision necessary to reach the same level of survey completeness for making a Super-Earth $M_p \sin i$ detection at P_{HZ} given the same amount of observing epochs as the transit survey. In this case the goal of the radial velocity survey is to make a sufficient number of $M_p \sin i$ detections in order to identify at least one that also transits. For the example at $0.2 M_\star$ the circular orbit semi-amplitude of a Super-Earth at P_{HZ} is $K \sim 5.2 \text{ m s}^{-1}$. To complete the analysis, we must relate n of a transit survey to an equivalent $n_{\text{obs},rv}$ for a radial velocity survey. The *MEarth* survey cycles between targets with a 5 min cadence on average and employs eight independent telescopes, and the *HARPS* M dwarf survey had a fixed exposure time of 15 min. Thus, for the Super-Earth survey comparison, we assume $n = X n_{\text{obs},rv}$, where $X = 24$, implying 24 observing epochs over independent targets are obtained in the transit survey for each radial velocity measurement. In the case of a transit survey with multiple targets per field, the conversion factor can be significantly larger (X is >4 orders of magnitude larger for a survey like *Kepler*; Borucki et al. 2010). Using the same completeness probability $p_c = 0.75$ for the radial velocity survey with a false alarm probability of $F = 0.001$, $n_{\text{obs},rv} = n/24$, and $m = n_{\text{obs},rv}$, we use Equation 7 to estimate the limiting semi-amplitude precision ratio, $K/\sigma_{rv} = 2.8$. Hence, for the $0.2 M_\star$ case, $\sigma_{rv} = 1.85 \text{ m s}^{-1}$. To summarize, a radial velocity survey will have reached the same survey completeness probability for a Super-Earth orbiting at the inner HZ for a late M dwarf as a simulated trigger-mode transit survey if it obtains $n_{\text{obs},rv}$ with a precision of $\sigma_{rv} = 1.85 \text{ m s}^{-1}$. If the radial velocity precision is $\sigma_{rv} > 1.85 \text{ m s}^{-1}$, then the radial velocity survey will not have reached the same level of survey completeness as the transit survey and is not as efficient for surveying $M_p \sin i$ detections that also transit their host.

For comparison, the mode of the radial velocity precision from the *HARPS* M dwarf survey sample is approximately equal to $\sigma_{\text{HARPS}} = 3.0 \text{ m s}^{-1}$ (Bonfils et al. 2011). In the above example of a Super-Earth at the inner HZ of $M_\star = 0.2 M_\odot$ host star, the inequality, $\sigma_{\text{HARPS}} \gtrsim \sigma_{rv}$, implies that formally the efficiency for transiting planet detection for *MEarth* is slightly greater than that for transiting planet detections among the *HARPS* M dwarf survey $M_p \sin i$ detections. The transiting planet yield not only depends upon relative efficiency, but also total observing time. The advantage for *MEarth* may even be larger given that the *MEarth* telescopes are dedicated to the survey whereas *HARPS* is a shared instrument.

The above results depend on the definition of the inner-HZ (Selsis et al. 2007; Kopparapu et al. 2013; Zsom et al. 2013). To first order, the results of Section 6.4 show that for a transit survey for fixed detection completeness $n \propto (P/\tau_{\text{dur}}) \propto P^{2/3}$. The expected scaling with P for a radial velocity survey maintaining the same detection completeness $n_{\text{obs},rv} \propto K^{-2} \propto P^{2/3}$ (see Equation 26 of Cumming (2004)) is the same. Numerical




FIG. 4.— Estimated radial velocity precision (σ_{rv}) needed to match the efficiency of the 8-telescope *MEarth* system for finding a transiting inner HZ Super-Earth (GJ 1214 analog) as a function of stellar mass with at least one, two, three, and five transit events required for detection (solid lines), from bottom to top, respectively. (Dashed line) shows results for the single transit triggering mode of detection in a transit survey, and the horizontal dotted line indicates the mode of the radial velocity precision achieved by the *HARPS* M dwarf survey (Bonfils et al. 2011). The calculation assumes $p_c = 0.75$ and $X = 24$.

calculations confirm this expected weak dependence on P . For the $0.2 M_\star$ case, $\sigma_{rv} = 1.9 \text{ m s}^{-1}$ and 1.7 m s^{-1} for a cooler, $T_{\text{eq}} = 240 \text{ K}$, and hotter, $T_{\text{eq}} = 300 \text{ K}$ inner HZ definition, respectively.

The discovery of GJ 1214 did not rely on the triggering mode for detection, but was detected after phasing two potential transit events (Charbonneau et al. 2009). Figure 4 summarizes the radial velocity precision required in order to reach the same level of completeness for a single target using the same amount of observing time for the transit survey design of requiring $k \geq 1, 2, 3$, and 5 events as the solid lines from bottom to top, respectively. The dash line shows results for the triggering mode of operation. The achieved precision of the *HARPS* M-dwarf survey $\sigma_{\text{HARPS}} = 3.0$ is also shown (horizontal dotted line) demarcates the stellar hosts where a 8-telescope system like *MEarth* is more (below) and less (above) efficient. In the triggering discovery mode (dashed line), $M_\star > 0.15 M_\odot$ is the stellar host regime where the 8-telescope *MEarth* system in the triggering discovery mode is more efficient than *HARPS* for detecting transiting Super-Earth planets. For a 3-transit requirement detection mode, the detection efficiency equivalency is at $M_\star \sim 0.22 M_\odot$.

The results shown in Figure 4 and the previous discussion assume the goal of the survey is to find a HZ planet that actually transits. Given the small probability to transit, the radial velocity survey will achieve a large number of detections $\propto 1/p_{\text{tr}}$ for every single HZ planet that transits. If the goal of the radial velocity survey is just to detect a single $M_p \sin i$ Super-Earth mass object, then it is more appropriate to compare to the time necessary for a transit survey to result in a single detection, $n_{\text{obs},tr}/p_{\text{tr}}$. The requisite radial velocity precision for a single radial velocity detection of a Super-Earth with the equivalent observing time for a single transiting planet detection using the 8-telescope *MEarth* system is




FIG. 5.— Same as Figure 4, but for a single $M_{\text{sin}} = 6.6 M_{\oplus}$ planet detection that does not necessarily transit.

shown in Figure 5. For measuring the occurrence rate of Super-Earth $M_p \text{ sin}i$ planets (Bonfils et al. 2011), the *HARPS* M dwarf survey is predicted to have superior efficiency for measuring Super-Earth population statistics compared to the *MEarth* survey. However, a transit survey observing fainter targets can achieve a much larger conversion factor, X . A transit survey requiring $k \geq 3$ transits for detection with a multi-plexing conversion factor of $X \sim 2200$ has equivalent detection efficiency to a radial velocity survey with $\sigma_{rv} = 2.0 \text{ m s}^{-1}$ for inner HZ Super-Earth planets around late M dwarfs.

One caveat to the discussion is the need for $1/p_{\text{tr}} > 56$ $M_p \text{ sin}i$ detections at the inner HZ for late M dwarfs before one expects to find one that also transits. Both the *HARPS* and *MEarth* surveys target the brightest M dwarfs available, but due to the geometric transit alignment one must survey a larger volume and fainter magnitudes for the $M_p \text{ sin}i$ detections that also transit. For uniform stellar density, the brightest transiting planet at the inner HZ for late M dwarfs will typically be $\Delta_{\text{mag}} \sim 3.0$ mag fainter than the brightest $M_p \text{ sin}i$ nontransiting detection. At the median brightness of the *MEarth* M dwarf sample ($V=15.2$, J. Irwin, *private communication*) the *HARPS* instrumental radial velocity precision is predicted to be $\sigma_{rv} \sim 10 \text{ m s}^{-1}$ (Bonfils et al. 2011). Thus at the median brightness of the *MEarth* M dwarf sample, *MEarth* is much more efficient at finding transiting Super-Earth planets at the inner HZ than if attempting to survey the *MEarth* M dwarf sample for transiting planets with *HARPS*.

4.2. Bright M Dwarfs Hosting A Transiting HZ Earth Analog

We repeat the analysis of the previous section down to the regime of truly Earth-size and Earth-mass planets at the inner HZ of the brightest M dwarfs. The *TESS* mission (Ricker et al. 2010) is one possibility for identifying transiting HZ Earth analogs orbiting bright stars, but here we briefly explore the possibility of a scaled-up *MEarth* survey and the associated equivalent radial velocity precision.

From the perspective of window functions, the required $n_{\text{obs}, tr}$ is the same as for the detection of Super-Earths at




FIG. 6.— Radial velocity precision needed to match the efficiency of a single-target transit survey for finding a transiting inner HZ Earth analog as a function of M_* where the transit survey requires at least one, one with triggering (dotted line), three, and five transit events for detection, bottom to top, respectively. The calculation assumes $p_c = 0.75$ and $X = 10$.

the inner HZ. However, the photometric precision must be improved by a factor ~ 7 from the Super-Earth survey to detect the shallower Earth-analog transit depth. In principle, the requisite photometric precision for enabling the trigger mode of detection for Earth-size planets orbiting M dwarfs has been demonstrated from the ground. Scintillation limited performance using 4m class telescopes has achieved $\sigma_p = 250 \text{ ppm min}^{-1}$ photometric noise rate on $V \sim 12$ for fields with a comparable brightness and spectral type companion in the optical (Gilliland et al. 1993; Burke 2010). Mann et al. (2011) achieve $1250 \text{ ppm min}^{-1}$ photometric noise rate on $V \sim 13$ late K and early M dwarfs with a 2m class telescope with an innovative target cycling approach. Thus, the practical limitation of a scaled up *MEarth* survey is available observing time on larger telescopes rather than a physical limit imposed by the Earth's atmosphere.

Carrying through the above calculation for an Earth analog at the inner HZ, we find $\sigma_{rv} = 0.58 \text{ m s}^{-1}$ is needed for comparable efficiency to the triggering mode transit survey. For this result we assume $X = 10$, which would be expected given the efficiency difference between a high resolution spectrograph for precision radial velocity measurements compared to single-target photometry. Since radial velocity instrumentation is typically on larger aperture telescopes¹⁰, assuming more efficient radial velocity observations, $X = 5$, results in $\sigma_{rv} = 0.87 \text{ ms}^{-1}$. The noise floor for M dwarf precision radial velocities was $\sigma_{rv} = 1.1 \text{ m s}^{-1}$ for the *HARPS* M dwarf survey (Bonfils et al. 2011). Results for additional transit survey designs of 2, 3, or 5 transits for detection are shown in Figure 6, and the precision needed to reach comparable efficiency of an $M_p \text{ sin}i$ detection is shown in Figure 7.

5. CONCLUSION

¹⁰ The multi-telescope radial velocity concept (Bottom et al. 2013) is another possibility – <http://www.astro.caltech.edu/minerva/Home.html>




FIG. 7.— Same as Figure 6, but for a single $M_{\text{sin}} = 1 M_{\oplus}$ planet in the inner HZ that does not necessarily transit. Note the change in vertical scale compared to Figure 6.

Pushing the ground-based photometric and radial velocity techniques provides an opportunity for detecting true Earth-sized planets at the inner HZ orbiting the brightest late M dwarfs. For example a scaled up *MEarth* survey could detect HZ Earth-sized transiting planets among the brightest late M dwarfs. We predict that for the same amount of observing time, a radial velocity survey requires precision $\sigma_{rv} < 0.6 \text{ m s}^{-1}$ in order to more efficiently detect a $M_p \text{ sin } i = 1 M_{\oplus}$ transiting planet at the inner HZ for mid M dwarf. From space, the *TESS* mission (Ricker et al. 2010) plans to survey early M and K dwarfs for such Earth analogs. There is great impetus for finding the rare Earth analogs that also transit among the brightest stars to take full advantage of the follow-up potential of *HST* and the upcoming opportunities provided by *JWST*. *JWST* will usher in an unprecedented capability for measuring the atmospheric constituents of potentially habitable planets (Valenti et al. 2005; Deming et al. 2009).

We demonstrate that the window function for a transit survey can be expressed in terms of the hypergeometric probability distribution. An analytic approximation to the window function simplifies the design of a transit survey and provides an accurate estimate to the observing time necessary to achieve a given level of completeness for detection of a transiting planet. Furthermore, to first order, the resulting window function for a transit survey depends only on the total number of independent observations and secondarily depends upon the distribution of observation times or baseline over which the observations are obtained. We employ these results regarding the transit survey window function model in order to determine the comparable radial velocity survey precision that achieves the same completeness level given the same number of observations. For the brightest M dwarfs, $V < 12$, we predict *HARPS* and the *MEarth* surveys have approximately the same efficiency for detecting Super-Earth planets that transit and orbit at the inner HZ. However, at the median brightness of the *MEarth* sample, $V = 15$, the *MEarth* survey is predicted to have much higher efficiency at detecting transiting Super-Earths than *HARPS* due to the

degraded radial velocity precision for fainter stars. The higher efficiency is predominately driven by the small geometric transit probability at the inner HZ for late M dwarfs. *HARPS* is predicted to have much higher efficiency in $M_p \text{ sin } i$ detections than *MEarth* has for transiting planets, thus making *HARPS* efficient at determining Super-Earth $M_p \text{ sin } i$ population statistics. However, complementary planet population studies are possible from wide field transit surveys that observe many targets per observing epoch ($X \sim 10^5$ in the case of *Kepler*; Youdin 2011; Howard et al. 2012; Dong & Zhu 2013; Fressin et al. 2013; Dressing & Charbonneau 2013; Petigura et al. 2013).

6. APPENDIX: ANALYTIC APPROXIMATIONS TO THE WINDOW FUNCTION

6.1. Binomial

Prior work (e.g. Eq. A2 of Deeg et al. 2004) adopted approach A of Section 2, “each time the planet transits, were we observing?” For this statistical model of the window function each orbital period time interval represents an independent trial. Specifically, the orbital period time interval represents a binomial trial with probability of our witnessing it equal to the average fraction of time spent observing f_{cov} . Thus, the probability of witnessing exactly k transits is

$$p_k = \binom{M}{k} f_{\text{cov}}^k (1 - f_{\text{cov}})^{M-k}, k = 0, 1, 2, \dots, M \quad (8)$$

recalling that the expectation value for the number of trials $M = T_{\text{obs}}/P$ (Eq. 2) and that the probability of witnessing k or more transits $p_{\geq k}$ is given by Eq. 3.

6.2. Poisson

Approach A also can be addressed, in appropriate limits, specifically large M and small f_{cov} (e.g. Appendix 4 of Lyons 1996), using the Poisson approximation to the binomial distribution,

$$p_k = \frac{\lambda^k}{k!} e^{-\lambda}, k = 0, 1, 2, \dots, M \quad (9)$$

where $\lambda = M f_{\text{cov}}$. Even though the Poisson distribution is defined for all $k \geq 0$, the maximum number of transits in time span T_{obs} is $M = \lfloor T_{\text{obs}}/P \rfloor + 1$. The physical interpretation of the Poisson process would be that transits are observed at random times at an average rate of λ transits in T_{obs} , which approaches f_{cov}/P in the appropriate limit of large M . Of course, transits do not occur at random times, but the approximation works if observations are sparsely and randomly distributed throughout the time span T_{obs} . Deeg et al. (2004) recommend that the Poisson approximation gives good results so long as $f_{\text{cov}} \lesssim 0.5$ and M is large. While it is the least accurate, the Poisson approximation is the simplest to compute; we recommend it for hand calculations (see Section 6.4) and to validate more complicated window function implementations.

6.3. Hypergeometric

In approach B, “each time we observe, is the planet transiting?”, the statistical model of the window function adopts the transit duration time interval as the relevant independent trial. We make an analogy between




FIG. 8.— Planetary transit observability can be modeled as drawing balls without replacement. A schematic of a transit light curve, with $P = 6$ units and with depth and width exaggerated for clarity, is above a set of colored balls. Each ball in the contiguous line of balls corresponds to a time interval equal to the transit duration. Between the first and last observations, there are $N = 12$ time intervals total: $n = 5$ intervals (solid circles) each with sufficient observational sensitivity to detect a transit if it had occurred during the given interval, and seven intervals (dashed circles) each with insufficient or no observations during the given interval. Red balls (labeled) correspond to $M = 2$ intervals in which a transit occurs; all other balls are white (no label).

searching for a transiting planet and randomly drawing colored balls from an urn without replacement (Figure 8). Each ball represents a time interval equal to the transit duration τ_{dur} and also a potential observation of a transit. The parameter n in the hypergeometric distribution is the number of unique observation intervals of length τ_{dur} that have sufficient sensitivity to detect a transit if it had occurred during the interval. Parceling the observing span T_{obs} into intervals of τ_{dur} results in a set of $N = T_{obs}/\tau_{dur}$ time intervals, or by analogy N balls in the urn. In the illustration, the orbital period P is six times the transit duration τ_{dur} . In general, for circular orbits and central transits, the ratio $P/\tau_{dur} \approx \pi a/R_*$, where a is the orbital semimajor axis and the formula neglects the planetary radius. The illustration shows a contiguous set of $N = T_{obs}/\tau_{dur} = 12$ balls, where $M = 2$ balls are red (in transit), and the $N - M = 10$ remaining balls are colored white (out-of-transit). There are $n = 5$ balls drawn with solid outlines (observations) and $N - n = 7$ balls with dashed outlines (observation gaps or observation intervals with insufficient sensitivity to detect a transit even if it had occurred in the particular interval). Thus the probability of having k observations occur during transit among the n observations is given by the hypergeometric distribution,

$$p_k = \frac{\binom{M}{k} \binom{N-M}{n-k}}{\binom{N}{n}}, k = 0, 1, 2, \dots, M \quad (10)$$

which is appropriate for this case of sampling without replacement. In the nomenclature of statistics,

$$M = \text{sub population size}, \quad (11)$$

$$N = \text{total population size, and} \quad (12)$$

$$n = \text{sample size.} \quad (13)$$

A limitation common to the binomial and hypergeo-

metric distributions is that their input parameters formally are required to be integers, because of the combinatorics operator “choose.” For example, each equation contains M choose k ; that k is an integer is not a problem, but if M is forced to also, then the window function can have a stair-step appearance. This is readily mitigated by using implementations of the operators that can handle real-valued inputs via the gamma function, e.g.

$$\binom{M}{k} = \frac{M!}{k!(M-k)!} = \frac{\Gamma(M+1)}{\Gamma(k+1)\Gamma(M-k+1)} = \frac{M}{k(M-k)} \frac{\Gamma(M)}{\Gamma(k)\Gamma(M-k)} \quad (14)$$

One may wish to use Stirling’s formula or a similar approximation for the factorials, or instead of the gamma function itself, one may wish to use a log(gamma) function if available, to avoid numerical overflow.

It is possible to show the numerical similarities of the binomial model of Deeg et al. (2004) with the hypergeometric model of this study by expressing $n = f_{cov}T_{obs}/\tau_{dur}$, then the binomial model can be written as $B(k; M, f_{cov}) = B(k; T_{obs}/P, n\tau_{dur}/T_{obs})$. The expectation of the binomial model of Deeg et al. (2004), $E[B(M, f_{cov})] = f_{cov}M = f_{cov}T_{obs}/P = n\tau_{dur}/P$, is the same as the expectation of the hypergeometric model of this study, $E[H(N, M, n)] = nM/N = n\tau_{dur}/P$. The approaches’ higher moments are not identical; For example, the variance ratio between the models, $Var[B(M, f_{cov})]/Var[H(N, M, n)] = (1 - \tau_{dur}/T_{obs})/(1 - \tau_{dur}/P)$. However, since $\tau_{dur} \ll P < T_{obs}$, for parameters generally applicable to a transit survey, the variances are nearly equal.

6.4. Probability of Witnessing at least one Transit or two Transits

The probability of witnessing at least one transit (from Equation 3) is

$$p_{\geq 1} = e^{-\lambda} \quad \text{Poisson, (15)}$$

$$= 1 - (1 - f_{cov})^M \quad \text{binomial, (16)}$$

$$= 1 - \frac{\binom{N-M}{n}}{\binom{N}{n}} \quad \text{hypergeometric, (17)}$$

Associating $f_{cov} = n\tau_{dur}/T_{obs}$, Equation 15 is readily inverted: $n = -\ln(1 - p_c)(P/\tau_{dur})$. This provides a convenient benchmark for planning purposes as well checking how probable a potential planetary discovery is in a transit search.

The probability of witnessing at least two transits is

$$p_{\geq 2} = e^{-\lambda}(1 - \lambda) \quad \text{Poisson, (18)}$$

$$= 1 - (1 - f_{cov})^M - Mf_{cov}(1 - f_{cov})^{M-1} \quad \text{binomial, or (19)}$$

$$= 1 - \frac{\binom{N-M}{n}}{\binom{N}{n}} - M \frac{\binom{N-M}{n-1}}{\binom{N}{n}} \quad \text{hypergeometric, (20)}$$

6.5. Comparisons to Numerical Simulations

In Figure 1 we compare the resulting window function given by Equation 3 to the exact numerical window function for a hypothetical ground-based transit survey that observes for 30 consecutive days with observations for 8 hours each night. The specified transit survey is the same as presented in Figure 1 of Beatty & Gaudi




FIG. 9.— Detection probability (upper solid black line) for a representative target observed by *Kepler* for 16 quarters and requiring $k \geq 3$ transits for detection. Detection probability (lower solid black line) for a *Kepler* target impacted by yearly data gaps of one missing quarter per year. Analytic window functions are shown using the hypergeometric approximation (Equations 3 and 10) for three or more transits, $k \geq 3$ (solid red lines). Also shown is the equivalent set of lines as above but requiring $k \geq 6$ transits for detection (dash lines) and the analytical window function for continuous observations with no gaps from Fleming et al. (2008) (blue solid line).

(2008) to demonstrate their analytical representation of a transit survey (solid blue line). The exact numerical window function (dash black line) shows the substantial alias structure that is typical of ground-based transit surveys due to the day-night observing cycle, and which is not modeled by the simpler analytic approximations.

The analytic expressions approximate well the average window function. To demonstrate this we show in Figure 1 a moving average of the exact numerical function using a 1 day window (solid black line). The left and right panels provide results for requiring $k \geq 2$ and $k \geq 3$ transit events for detection, respectively. We show the hypergeometric window function model from this study (Equation 10) as the solid red line. Solar values for the stellar parameters are assumed. The Poisson window function model (Equation 9, dash red line) also does a reasonable job in this scenario with $f_{\text{cov}} = 0.33$.

Figure 3 shows the window function for a transit survey that operates in a low duty cycle mode obtaining several observations per night over the course of an observing season. We simulate this mode of operation through specifying a time series with $n = 200$ that is expected for a target from a single observatory over the course of an observing season including losses due to weather and stellar parameters appropriate for a mid M dwarf star. The exact numerical window functions assuming at least $k \geq 1, 2, 3$, and 5 transit events are observed (solid lines from top to bottom, respectively). The hypergeometric model (solid red line) and its Poisson model approximation (dash blue line) agree extremely well as expected for a low duty cycle transit survey.

Figure 9 shows the window function for two typical targets observed by *Kepler* using Quarters 1 through 16 (~ 3.6 yr) of nearly continuous coverage

(Tenenbaum et al. 2013). The rightmost set of black solid, red solid, and blue solid lines correspond to the coverage of *Kepler*-11, host to six transiting planets (Lissauer et al. 2011). This target was observed for all 16 *Kepler* Quarters. For the time series, we employ barycentric julian date mid-cadence times stamps that have valid data as determined by the *Kepler* pipeline (Smith et al. 2012; Wu et al. 2010). The hypergeometric model (solid red line) requiring $k \geq 3$ transit events being observed agrees well with the exact numerical calculation (solid black line) also requiring $k \geq 3$. As expected the analytical window model function for continuous observations with no gaps of (Table 2 of Fleming et al. 2008), overestimates the window function due to the reality of gaps in *Kepler* data. The leftmost set of solid black and solid red lines correspond to the coverage of the host to *TrES*-2b (a.k.a. *Kepler*-1b; O’Donovan et al. 2006b; Barclay et al. 2012). This target is impacted by the loss of a detector in Quarter four (Batalha et al. 2013), causing missing data every fourth *Kepler* Quarter. The detector loss impacts $\sim 19\%$ of *Kepler* targets having reduced amount of data available. The dash lines are the same as above, but they require $k \geq 6$ transits for detection.

The hypergeometric model does not fully capture the details present in the exact numerical window functions. The assumptions of the hypergeometric model begin to break down in observing scenarios consisting of high duty cycle blocks interspersed with long periods of no data. In addition to the sharp, isolated aliases at the diurnal cycle and its harmonics prevalent in ground-based transit survey window functions (which we readily average over), there are “broad-band” alias structures present across all orbital periods that can result in the exact numerical window function to deviate from the hypergeometric model.

In Figure 2 we demonstrate the principal dependence of a transit survey window function on n and the weak dependence on T_{obs} . We show the hypergeometric model (solid lines) and the moving average of the exact numerical window function (dash lines) for three scenarios that have the same n but vary in T_{obs} . The left and right panels vary by requiring $k \geq 2$ and 3 transit events for detection, respectively. The baseline transit survey consists of 7 hours of observations each day for 30 consecutive days (black lines). We simulate a longitudinally-distributed network survey by considering a transit survey with half T_{obs} (15 consecutive days) of the baseline case while doubling the observing time each day to conserve n (red lines). The window function does show an increase in detection probability at intermediate periods at the cost of removing the low detection probability tail toward longer periods of the baseline scenario. The third n -conserving scenario consists of splitting the 30 consecutive days of observations from the baseline case into two 15 consecutive day groups separated by a year (blue lines) giving a $T_{\text{obs}} = 380$ day. In contrast to the n -conserving cases, we show a transit survey window function resulting from doubling n while keeping T_{obs} the same as the baseline case (magenta lines). These scenarios demonstrate that the detection probability of a transit survey are very sensitive to n .

This work benefited from discussions with Cullen

TABLE 1
Stellar, Transit, & Radial Velocity Parameters

Blake, B. Scott Gaudi, Jonathan Irwin, Jon Jenkins, and George Ricker. We thank the referee for insightful suggestions which improved the manuscript.

M_* [$M_\odot$]	R_* [$R_\odot$]	T_{eff} [K]	P_{HZ} [day]	p_{tr}	$\Delta_\oplus$	$\Delta_{\text{S}\oplus}$ [ppm]	τ_{dur} [ppm]	n^a [hr]	n_{ngt}	n/p_{tr}	$K_{\text{S}\oplus}$ [m s $^{-1}$]	$\sigma_{\text{rv S}\oplus}^b$ [m s $^{-1}$]	$K_\oplus$ [m s $^{-1}$]	$\sigma_{\text{rv}\oplus}^c$ [m s $^{-1}$]
0.15	0.20	3077	10.47	0.0188	2167	15801	1.18	492	73	26139	6.83	2.69	1.035	0.81
0.17	0.22	3109	11.62	0.0184	1724	12568	1.29	472	76	25606	6.07	2.29	0.920	0.70
d 0.20	0.26	3155	13.61	0.0179	1266	9233	1.46	449	82	25086	5.17	1.85	0.783	0.58
0.22	0.28	3185	15.13	0.0176	1052	7671	1.59	436	87	24825	4.68	1.62	0.709	0.51
0.25	0.32	3230	17.75	0.0171	818	5967	1.82	421	96	24655	4.08	1.35	0.618	0.44
0.27	0.35	3261	19.75	0.0168	703	5127	1.99	413	102	24653	3.74	1.21	0.566	0.40
0.30	0.38	3310	23.14	0.0163	572	4171	2.26	404	114	24849	3.30	1.04	0.501	0.34
0.35	0.44	3404	29.86	0.0154	426	3110	2.75	404	139	26278	2.74	0.86	0.415	0.28
0.40	0.50	3520	37.76	0.0144	336	2449	3.26	418	170	29071	2.32	0.76	0.351	0.25
0.45	0.55	3661	46.32	0.0133	278	2025	3.69	442	204	33257	2.00	0.70	0.303	0.22
0.50	0.59	3811	55.12	0.0123	239	1745	4.06	473	240	38559	1.76	0.67	0.267	0.20
0.55	0.63	3952	64.07	0.0114	213	1555	4.38	503	276	44098	1.57	0.63	0.238	0.19
0.60	0.66	4105	74.87	0.0106	192	1402	4.75	538	319	50889	1.41	0.61	0.213	0.18
0.70	0.75	4591	117.39	0.0085	148	1078	5.95	657	489	77718	1.09	0.56	0.166	0.15
0.80	0.86	5198	189.37	0.0066	114	834	7.49	827	775	125410	0.85	0.53	0.129	0.14
0.90	0.90	5509	231.34	0.0059	103	749	8.15	924	941	157421	0.73	0.49	0.112	0.13
1.00	1.03	5794	307.04	0.0053	80	581	9.78	1011	1235	190520	0.63	0.44	0.095	0.11
1.10	1.16	6063	404.58	0.0048	63	459	11.76	1097	1613	226371	0.54	0.40	0.081	0.10

^a Single transit triggering mode of operation and $p_c = 0.75$ completeness level.

^b Assumes $X = 24$ (see Section 4.1).

^c Assumes $X = 10$ (see Section 4.2).

^d The bold face row is discussed in Section 4.1.

REFERENCES

- Alonso, R., Brown, T. M., Torres, G., et al. 2004, *ApJ*, 613, L153
 Alsubai, K. A., Parley, N. R., Bramich, D. M., et al. 2011, *MNRAS*, 417, 709
 Bakos, G. Á., Noyes, R. W., Kovács, G., et al. 2007, *ApJ*, 656, 552
 Barclay, T., Huber, D., Rowe, J. F., et al. 2012, *ApJ*, 761, 53
 Barclay, T., Burke, C. J., Howell, S. B., et al. 2013, *ApJ*, 768, 101
 Barnes, J. W. 2007, *PASP*, 119, 986
 Batalha, N. M., Borucki, W. J., Bryson, S. T., et al. 2011, *ApJ*, 729, 27
 Batalha, N. M., Rowe, J. F., Bryson, S. T., et al. 2013, *ApJS*, 204, 24
 Beatty, T. G., & Gaudi, B. S. 2008, *ApJ*, 686, 1302
 Beatty, T. G., Pepper, J., Siverd, R. J., et al. 2012, *ApJ*, 756, L39
 Berta, Z. K., Charbonneau, D., Bean, J., et al. 2011, *ApJ*, 736, 12
 Berta, Z. K., Irwin, J., & Charbonneau, D. 2013, *ApJ*, 775, 91
 Boisse, I., Bouchy, F., Hébrard, G., et al. 2011, *A&A*, 528, A4
 Bonfils, X., Delfosse, X., Udry, S., et al. 2011, submitted *A&A*, arXiv:1111.5019
 Borucki, W. J., Koch, D., Basri, G., et al. 2010, *Science*, 327, 977
 Brown, T. M. 2003, *ApJ*, 593, L125
 Burke, C. J., Gaudi, B. S., DePoy, D. L., & Pogge, R. W. 2006, *AJ*, 132, 210
 Burke, C. J. 2008, *ApJ*, 679, 1566
 Burke, C. J. 2010, in *Towards other Earths: Perspectives and limitations in the ELT era*, ed. N. C. Santos, C. Melo, L. Pasquini, A. Glindemann (Porto: Centro de Astrofísica da Universidade do Porto), Session 1, http://www.astro.up.pt/investigacao/conferencias/toe2009/DVD/TOE/TOE_mackinnon.pdf
 Bottom, M., Muirhead, P. S., Asher Johnson, J., & Blake, C. H. 2013, *PASP*, 125, 240
 Charbonneau, D., Brown, T. M., Latham, D. W., & Mayor, M. 2000, *ApJ*, 529, L45
 Charbonneau, D., Berta, Z. K., Irwin, J., et al. 2009, *Nature*, 462, 891
 Collier Cameron, A., Bouchy, F., Hébrard, G., et al. 2007, *MNRAS*, 375, 951
 Cosentino, R., Lovis, C., Pepe, F., et al. 2012, *Proc. SPIE*, 8446
 Covino, E., Esposito, M., Barbieri, M., et al. 2013, *A&A*, 554, A28
 Cox, A. N. 2000, *Allen's Astrophysical Quantities*
 Cumming, A. 2004, *MNRAS*, 354, 1165
 Deeg, H. J., Alonso, R., Belmonte, J. A., et al. 2004, *PASP*, 116, 985
 Deming, D., Seager, S., Winn, J., et al. 2009, *PASP*, 121, 952
 Dong, S., & Zhu, Z. 2013, *ApJ*, 778, 53
 Dressing, C. D., & Charbonneau, D. 2013, *ApJ*, 767, 95
 Fleming, S. W., Kane, S. R., McCullough, P. R., & Chromey, F. R. 2008, *MNRAS*, 386, 1503
 Fressin, F., Torres, G., Désert, J.-M., et al. 2011, *ApJS*, 197, 5
 Fressin, F., Torres, G., Charbonneau, D., et al. 2013, *ApJ*, 766, 81
 Gaudi, B. S. 2000, *ApJ*, 539, L59
 Gaudi, B. S., Seager, S., & Mallen-Ornelas, G. 2005, *ApJ*, 623, 472
 Gillon, M., Pont, F., Demory, B.-O., et al. 2007, *A&A*, 472, L13
 Gilliland, R. L., Brown, T. M., Kjeldsen, H., et al. 1993, *AJ*, 106, 2441
 Gilliland, R. L., Brown, T. M., Guhathakurta, P., et al. 2000, *ApJ*, 545, L47
 Gomes da Silva, J., Santos, N. C., Bonfils, X., et al. 2012, *A&A*, 541, A9
 Henry, G. W., Marcy, G. W., Butler, R. P., & Vogt, S. S. 2000, *ApJ*, 529, L41
 Horne, J. H., & Baliunas, S. L. 1986, *ApJ*, 302, 757
 Howard, A. W., Marcy, G. W., Bryson, S. T., et al. 2012, *ApJS*, 201, 15
 Irwin, J. M., Quinn, S. N., Berta, Z. K., et al. 2011, *ApJ*, 742, 123
 Kane, S. R., Mahadevan, S., von Braun, K., Laughlin, G., & Ciardi, D. R. 2009, *PASP*, 121, 1386
 Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, *Icarus*, 101, 108
 Kopparapu, R. K., Ramirez, R., Kasting, J. F., et al. 2013, *ApJ*, 765, 131
 Koppelman, M., Torres, G., Jha, S., & Sasselov, D. D. 2003, *Nature*, 421, 507
 Lissauer, J. J., Fabrycky, D. C., Ford, E. B., et al. 2011, *Nature*, 470, 53
 Mann, A. W., Gaidos, E., & Aldering, G. 2011, *PASP*, 123, 1273
 McCullough, P. R., Stys, J. E., Valenti, J. A., et al. 2006, *ApJ*, 648, 1228
 Nutzman, P., & Charbonneau, D. 2008, *PASP*, 120, 317
 O'Donovan, F. T., Charbonneau, D., Torres, G., et al. 2006, *ApJ*, 644, 1237
 O'Donovan, F. T., Charbonneau, D., Mandushev, G., et al. 2006, *ApJ*, 651, L61
 Petigura, E. A., Howard, A. W., & Marcy, G. W. 2013, *Proceedings of the National Academy of Science*, 110, 19273
 Queloz, D., Bouchy, F., Moutou, C., et al. 2009, *A&A*, 506, 303
 Ricker, G. R., Latham, D. W., Vanderspek, R. K., et al. 2010, *Bulletin of the American Astronomical Society*, 42, #450.06

- Saar, S. H., & Donahue, R. A. 1997, *ApJ*, 485, 319
- Seager, S., & Mallén-Ornelas, G. 2003, *ApJ*, 585, 1038
- Selsis, F., Kasting, J. F., Levrard, B., et al. 2007, *A&A*, 476, 1373
- Smith, J. C., Stumpe, M. C., Van Cleve, J. E., et al. 2012, *PASP*, 124, 1000
- Stevens, D. J., & Gaudi, B. S. 2013, *PASP*, 125, 933
- Stevenson, K. B., Harrington, J., Lust, N. B., et al. 2012, *ApJ*, 755, 9
- Tenenbaum, P., Jenkins, J. M., Seader, S., et al. 2013, *arXiv:1311.0248*
- Torres, G., Konacki, M., Sasselov, D. D., & Jha, S. 2004, *ApJ*, 614, 979
- Torres, G., Fressin, F., Batalha, N. M., et al. 2011, *ApJ*, 727, 24
- Valenti, J. A., Arribas, S., Ferruit, P., et al. 2005, *Bulletin of the American Astronomical Society*, 37, #115.05
- von Braun, K., Kane, S. R., & Ciardi, D. R. 2009, *ApJ*, 702, 779
- Winn, J. N., Matthews, J. M., Dawson, R. I., et al. 2011, *ApJ*, 737, L18
- Wolfe, D. E. & Gutman, S. I. 2000, *J. Atmos. Ocean. Technol.*, 17, 426
- Wu, H., Twicken, J. D., Tenenbaum, P., et al. 2010, *Proc. SPIE*, 7740, 774019
- Youdin, A. N. 2011, *ApJ*, 742, 38
- Zsom, A., Seager, S., de Wit, J., & Stamenković, V. 2013, *ApJ*, 778, 109